

Charge Density Distributions in Doped Antiferromagnetic Insulators

L. S. Isaev, A. P. Protogenov*

*Nizhny Novgorod State University, 603950 Nizhny Novgorod
Institute of Applied Physics of the RAS, 603950 Nizhny Novgorod*

Abstract

We consider the form of the charge density nano-scale configurations in underdoped states of planar antiferromagnetic insulators in the framework of a soft variant of Faddeev-Niemi model. It is shown that there is such a level of doping and the temperature range, where charge density distributions in the form of closed quasi-one-dimensional structures are more preferable.

PACS: 74.25.-q, 74.80.-g, 71.10.Hf, 71.10.-w

Low-dimensional inhomogeneities in the distribution of charge [1, 2, 3] and spin [4, 5] degrees of freedom at a moderate level of doping of planar antiferromagnetic insulators are the subject of active experimental and theoretical study nowadays. One of the reasons of increased attention to coherent quasi-one-dimensional charge structures is their existence in the state, which precedes the high-temperature superconducting phase. Keeping in mind this key property [6], for describing the mentioned phases we should choose such a model, which will contain them as limiting cases. In the framework of the mean field theory, the two-component Ginzburg-Landau model, defined on the complex projective space CP^1 , satisfies the searched requirements. It was shown recently [7] that there is its exact mapping to the following extended variant of the $\mathbf{n}$ -field model:

$$F = \int d^3x \left[\frac{1}{4} \rho^2 (\partial_k \mathbf{n})^2 + (\partial_k \rho)^2 + \frac{1}{16} \rho^2 \mathbf{c}^2 + (F_{ik} - H_{ik})^2 + V(\rho, n_3) \right]. \quad (1)$$

The free energy in Eq.(1) is defined by the density ρ^2 , as well as by the unit vector $\mathbf{n}$ and the momentum $\mathbf{c} = \mathbf{J}/\rho^2 = 2(\mathbf{j} - 4\mathbf{A})$. They characterize the spatial distributions of spin degrees of freedom with current $\mathbf{J}$ or without it. The total current contains the paramagnetic part $\mathbf{j} = i[\chi_1 \nabla \chi_1^* - c.c. + (1 \rightarrow 2)]$ and the diamagnetic term $-4\mathbf{A}$. The following notations were used in Eq.(3): $F_{ik} = \partial_i c_k - \partial_k c_i$ and $H_{ik} = \mathbf{n} \cdot [\partial_i \mathbf{n} \times \partial_k \mathbf{n}] \equiv \partial_i a_k - \partial_k a_i$. The vector a_i has the sense of the gauge potential which parametrizes the strength of the internal magnetic field. Provided the electron spin and charge are transferred from one of the four sites of some plackets to the dopant reservoir, the terms with H_{ik} in Eq. (1) characterize (in the infrared limit) the mean degree of non-collinearity $< 0 | \mathbf{S}_1 \cdot [\mathbf{S}_2 \times \mathbf{S}_3] | 0 >$ in the orientation of three spins remaining in the sites of a quadratic

*e-mail: alprot@appl.sci.nnov.ru

lattice placket. Therewith the deficit of the charge density ρ_h^2 relates to the density ρ^2 , which describes the distributions of the exchange integral in (1), as $\rho^2 + \rho_h^2 = \text{const}$. From the long-wavelength point of view the distribution of the spin density ρ^2 in the limited region with the exponential law of decrease at its boundary (for example, at the distribution in the circle of the radius r_0 with the exponential decrease at the length $R \ll r_0$) will be accompanied by the quasi-one-dimensional distribution of the charge density ρ_h^2 along the boundary of this region, i.e. along the ring with thickness R and radius r_0 . It is seen from here that studying spatial configurations of the density field ρ^2 in planar systems allows us to find the form of one-dimensional distributions of the electric charge density with the aid of the above-mentioned holographic projection.

The multipliers of the first term in the soft variant of the $\mathbf{n}$ -field model (1) describe the distributions of conjugated variables – the spin stiffness ρ^2 and the square of the inverse length $(\partial_k \mathbf{n})^2$ of the density screening. It follows from this particular example already, that the competition of the order parameters ρ , $\mathbf{n}$ and $\mathbf{c}$ may be the origin of the existence of phase states with various ordering of strongly correlated charge and spin degrees of freedom. We enumerate the main limiting cases of model (1) in inhomogeneous ($\mathbf{n} \neq \text{const}$) phase states as follows: (i) $\mathbf{c} = 0$, $\rho = \text{const}$; (ii) $\mathbf{c} = 0$, $\rho \neq \text{const}$; (iii) $\mathbf{c} \neq 0$, $\rho = \text{const}$; (iv) $\mathbf{c} \neq 0$, $\rho \neq \text{const}$. In the case of $\mathbf{n} = \text{const}$ and $\mathbf{c} \neq 0$, $\rho \neq \text{const}$ the functional (1) is equivalent to the one-component Ginzburg-Landau model.

The model for studying inhomogeneous distributions of spin degrees of freedom in the form of knots for case (i) was proposed in paper [8]. In this rigid phase $F \geq 32\pi^2 |Q|^{3/4}$ [9, 10], where $Q = 1/(16\pi^2) \int d^3x \varepsilon_{ikl} a_i \partial_k a_l$ is Hopf invariant. The inhomogeneous superconducting state (iii) with finite momentum of pairs and universal character of correlations was recently considered in [11]. In this phase the gain of the free energy $F \geq 32\pi^2 |Q|^{3/4} (1 - |L|/|Q|)^2$ depends on the index $L = 1/(16\pi^2) \int d^3x \varepsilon_{ikl} c_i \partial_k a_l$ of mutual linking degree of one-dimensional configurations, for which the fields $\mathbf{c}$ and $\mathbf{a}$ are defined.

The aim of this paper is to study the properties of states outside the superconducting phase from the second line (ii) of the list of limiting cases, to which the model (1) leads. In this soft version of the Faddeev-Niemi model [12, 13] the functional (1) has the following form

$$F = \int d^3x \left[\frac{1}{4} \rho^2 (\partial_k \mathbf{n})^2 + H_{ik}^2 + (\partial_k \rho)^2 - b\rho^2 + \frac{d}{2} \rho^4 \right]. \quad (2)$$

The state with the broken antiferromagnetic order has a lower energy than the "soft" state in which we are interested now. The last one may be metastable [14]. We will consider just such states and compare their contribution to the Ginzburg-Landau energy without studying the problems of their relaxation, the critical sizes of nuclei of different phases and etc., which are of separate interest.

It has been known for a long time (see, for example, [15]) that in such a phase state the distributions of the charge density ρ_h^2 have the form of stripes¹. It is seen from Eq.(2) that the loss in the free energy due to inhomogeneity may be reduced by developing inhomogeneous configurations of the field ρ with one-dimensional distribution of the gradient $\partial_k \rho$. From this point of view it is almost obvious that the quasi-one-dimensional field configurations ρ in the form of rings give the smallest contribution to the energy.

¹We suppose that the stripe characteristic sizes are essentially greater than the lattice scale. In this case, the use of phenomenological approach of the Ginzburg-Landau mean field theory is justified.

Whether these spin structures will be two-dimensional with sharp boundaries or one-dimensional and open, forming stripes, or closed almost one-dimensional structures in the form of rings, depends also on the parameters of the potential $V(\rho, n_3)$. In the phase $n_3 = \text{const}$ with neutral spin currents the answer will depend on the sign of the coefficient b in the potential $V(\rho, n_3)$. The positive constant b corresponds to the phase with the broken antiferromagnetic order in (2). In this case the form of quasi-one-dimensional structures will depend on the result of the competition between the first and the fourth terms in Eq. (2) (see also [16]). We will show that in this phase, far from T_c charge structures with the open ends are preferable, while in the case $T \rightarrow T_c$ we should prefer rings. The first experimental indication of the existence of charge structures in the form of rings in the underdoped phase of planar systems is in the recent paper [17]. In a certain sense, the tunnel microscope in this experiment [17] collects data from a two-dimensional slice of knots [12].

Let us find the contribution to the free energy (2) from quasi-one-dimensional distributions of the density ρ^2 in the form of a ring and a stripe and compare the results of the calculations with the experimental data. We choose the following trial functions for the configurations of the field ρ in the form of a ring and a stripe:

$$\rho = \rho_0 e^{-(r-r_0)^2/2R^2}, \quad (3)$$

$$\rho = \rho_0 \exp[-x^2/2L_x^2] \times \begin{cases} 1, & |y| \leq L_y, \\ \exp[-(|y| - L_y)^2/2L_x^2], & |y| > L_y. \end{cases} \quad (4)$$

Here $\rho_0 = \sqrt{b/d}$, r_0 is the ring radius, R is its width, $2L_y = 2\pi r_0$ is the stripe length, $L_x = R$ is its width. Since configurations (3) and (4) do not depend on the third coordinate, we shall assume that along this coordinate the size is limited by the length L_z , and also that $R < r_0$. The calculation of the energy (2) with the aid of (3) and (4) yields the following result for the contribution to the free energy from the spin degrees of freedom distributed over the ring F_r and the stripe F_{xy} :

$$F_r = \pi \rho_0^2 L_z \frac{\bar{r}_0}{R} \left(1 + \frac{R^2}{\xi^2} \right), \quad (5)$$

$$F_{xy} = \pi \rho_0^2 L_z \frac{\bar{r}_0}{R} \left(1 + \frac{R^2}{\xi^2} + \frac{R}{\bar{r}_0} + \left(n_0 - \frac{3}{4}b \right) \frac{R^3}{\bar{r}_0} \right). \quad (6)$$

Here $\bar{r}_0 = \sqrt{\pi} r_0$, $1/\xi^2 = 2[n_0 - (1 - 1/\sqrt{8})b]$, n_0 is a certain characteristic value of the "multiplier" $(\partial_k \mathbf{n})^2$ in (2), which is of the order $c_1 R^{-2}$, whereas $b = c_2 R^{-2} \delta T$, where $c_i \sim 1$, $\delta T = (T_c - T)/T_c$. In these equations we omitted the term H_{ik}^2 from Eq. (2) since we consider that it is the same for the both types of distributions.

At $R \ll r_0$ the optimal width of the charge structures $R = \xi$ and the difference of free energies $\Delta F = F_{xy} - F_r$ in units $\pi \rho_0^2 L_z$ equals $\Delta F = 1 + c_1 - (3/4)c_2 \delta T$. It follows from this equation that in the case of $4(1 + c_1)/3c_2 < 1$ in the temperature range $[1 - 4(1 + c_1)/3c_2] T_c < T < T_c$, which is contiguous with the critical temperature T_c of the transition to the state of the spin pseudo-gap, the contribution to the free energy from the rings is smaller than from the stripes (see Fig. 1). In the range of temperatures $T < T_c [1 - 4(1 + c_1)/3c_2]$ the stripes are the main configurations. As is known, one may reach T_c , keeping the temperature constant, by increasing the level of doping. Under




Figure 1: Schematic representation of closed (b) and open (c) quasi-one-dimensional structures of the charge density (see. [17]) around antiferromagnetic dielectric nano-clusters (a).

these experimental conditions studied in the recent paper [17], it was reported for the first time that ring-shaped charge structures in underdoped phase states were observed.

Let us make some remarks concerning the distribution of the spin density ρ^2 in the disk, surrounded by the ring charge distribution. All results were obtained under the assumption that the spin disorder arose only in the region directly adjacent to the disk edge. Therefore, there was the antiferromagnetic phase inside it. If the antiferromagnetic order is broken everywhere in the disk, it is necessary to consider the corresponding distribution of the density ρ^2 in the form of a disk with further comparison of its contribution with F_r . When considering the contribution of the density distributions ρ^2 to the free energy in the form of a disk, we get a double gain due to the existence only one edge and have a loss due to the area. The calculation shows that at small R/r_0 the distributions in the form of rings appear to be more preferable.

Let us consider the dependence of the critical temperature T_c on the level of doping. With this aim, we use the relation of F_r to F_{xy} in the following form:

$$\frac{F_r}{F_{xy}} = \frac{1}{1 + B R/\bar{r}_0}, \quad (7)$$

where

$$B = \frac{1}{2} \left[1 + \frac{1}{2} \frac{n_0 - 3/4b}{n_0 - 1/\sqrt{8}b} \right] = \frac{3}{4} \frac{n_0 - 0.68b}{n_0 - 0.65b}.$$

One can see that configurations in the form of stripes are more preferable in the range $0.65 < n_0/b < 0.68$ where $F_{xy} < F_r$. Normalizing the density ρ_0^2 to the particle number N by the condition $N = 2\pi\rho_0^2\xi L_z r_0 m$, we get the relation between the parameters n_0

and b in the form $x = b/\sqrt{n_0 - \alpha b}$. Here $x = Nd/\sqrt{2m\pi L_z r_0}$ and $\alpha = 0.65$. Thus, for the bounds of the above-considered range where $n_0 \sim b$, we have $T(x) = T_c(1 - Ax^2)$ with a certain constant A . Therefore, inside the region belonging to the phase state with the broken antiferromagnetic order, there exists a narrower region, located between the parabolas $T(x)$, where the charge structures have the form of stripes.

The analysis of the inhomogeneous superconducting state $F_{ik} \neq 0$, $\rho \neq \text{const}$ is an open problem at present. Since $(\partial_k \rho)^2 \neq 0$, the additional contribution decreases the gain in the free energy [11] in the superconducting phase. The superconducting state from this point of view will be studied more thoroughly in a separate paper. Here we mention only that the superconducting current with the amplitude $\mathbf{c}_0$, flowing along the rings (3), yields the term of the order $\mathbf{c}_0^2 R^2$ in the multiplier $(1 + R^2/\xi^2)$ of Eq. (5). This explains in particular why the superconducting region on the phase diagram "temperature - doping level" is shifted to the line $\delta T(x) = 0$ of the transition to the state with the spin pseudo-gap. Indeed, in this case $V_{eff}(\rho, n_3) = -b_{eff}\rho^2 + (d/2)\rho^4$ with $b_{eff} = b - (n_0 + \mathbf{c}_0^2) = (\text{const}/R^2)\delta T$. We see that the finite value of the momentum $\mathbf{c}$ of superconducting pairs decreases δT .

In conclusion, we have compared the contributions to the free energy from the charge density distributions with open and closed ends and found the regions of the existence of both types of configurations. The obtained results support the conjecture of the paper [16] and explains the results of the experiment described in [17].

We are grateful to S. Davis, L.D. Faddeev, S.M. Girvin, E.A. Kuznetsov, B. Lake, A.I. Larkin, A.G. Litvak, C. Renner, H. Takagi, V.A. Verbus, G.E. Volovik, Y.-S. Wu for advices and useful discussions. This paper was supported in part by the grant of the Russian Foundation for Basic Research No. 01-02-17225.

References

- [1] S. H. Pan *et al.*, Nature **413**, 282 (2001).
- [2] J. E. Hoffman *et al.*, Science **295**, 466 (2002).
- [3] S.J.L. Billinge, E.S. Bozin, M. Gutmann, and H. Takagi, "Microscopic charge inhomogeneities in underdoped $La_{2-x}Sr_xCuO_4$: local structural evidence", cond-mat/0005032.
- [4] B. Lake *et al.*, Nature **415**, 299 (2002).
- [5] B. Lake *et al.*, Science **291**, 1759 (2001).
- [6] S. Caprara, C. Castellani, C. Di Castro, M. Grilli, A. Perali, "Charge and spin inhomogeneity as a key to the physics of the high T_c cuprates", cond-mat/9907265.
- [7] E. Babaev, L. D. Faddeev, A. J. Niemi, Phys. Rev. B **65**, 100512 (2002).
- [8] L. D. Faddeev, "Quantization of solitons", Princeton report No. IAS-75-QS70, 1975.
- [9] A. F. Vakulenko, L. V. Kapitansky, Sov. Phys. Dokl. **24**, 433 (1979).
- [10] R. S. Ward, Nonlinearity **12**, 1 (1999), hep-th/9811176.

- [11] A. P. Protogenov, V. A. Verbus, Pis'ma v ZheTF, **76**, 60 (2002).
- [12] L. D. Faddeev, A. J. Niemi, Nature **387**, 58 (1997).
- [13] M. Lübke, S. M. Nasir, A. Niemi and K. Torokoff, Phys. Lett. **B534** 195 (2002).
- [14] A. Z. Patashinsky, B. I. Shumilo, Sov. ZhETP, **50**, 712 (1979).
- [15] C. Di Castro, L. Benfatto, S. Caprara, C. Castellani, M. Grilli, Phys. C **341-348**, 1715 (2000)
- [16] A. J. Niemi, "Phases of bosonic strings and two dimensional gauge theories", hep-th/0206227.
- [17] H. Takagi. *Real k-space imaging of electronic states in Na-doped $Ca_2CuO_2Cl_2$* . Talk presented at the Workshop on "Emergent materials and highly correlated electrons", Trieste, August 2002;
<http://agenda.ictp.trieste.it/agenda/current/fullAgenda.php?email=0&ida=a01152>